

Quantum theory of condensed matter I

PD Dr. Andrea Donarini

Tue 10:00 - 12:00 H33


Dr. Ivan Dmitriev

Thu 10:00 - 12:00 H33

Tue 12:00 - 14:00 9.2.01

Sheet 4

1. Electronic structure of non-interacting graphene



A graphene sheet is a honeycomb lattice of carbon atoms, as shown above. Let the distance between carbon atoms be $a_C = 1.42 \text{ \AA}$. A good model for graphene is to consider a plane in which there is one valence electron per carbon atom. To calculate the electronic bands of the non-interacting problem we will use the LCAO method with a single p_z orbital at each carbon site. Let \vec{R} denote the centers of the hexagons in the honeycomb: these form the underlying Bravais lattice. (Note that it is different from the graphene hexagonal lattice.) The unit cell spanned by \vec{a}_1 and \vec{a}_2 contains two carbon atoms conventionally labelled as A and B atoms, located at $\vec{R} + \vec{\tau}_A$, $\vec{R} + \vec{\tau}_B$ as shown in the figure. The tight-binding amplitude connecting these sites is denoted by $-t$. We can choose the zero of energy so that the energy of an isolated p_z orbital is zero. Then the entire Hamiltonian consists only of the nearest-neighbour hopping.

- Find the reciprocal lattice and construct the first Brillouin zone. (1 Point)
- Write the Hamiltonian in the basis of the extended states

$$\begin{aligned}
 |\phi_{A\vec{k}}\rangle &= \frac{1}{\sqrt{N_{cell}}} \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} |p_z, \vec{R} + \vec{\tau}_A\rangle \\
 |\phi_{B\vec{k}}\rangle &= \frac{1}{\sqrt{N_{cell}}} \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} |p_z, \vec{R} + \vec{\tau}_B\rangle,
 \end{aligned}$$

where N_{cell} is the number of the unit cells of the crystal and $|p_z, \vec{S}\rangle$ is the state of the p_z orbital localized in the position \vec{S} . Hint: make use of the relation

$$\frac{1}{N_{cell}} \sum_{\vec{R}} \exp\{i(\vec{k} - \vec{k}') \cdot \vec{R}\} = \delta_{\vec{k}\vec{k}'},$$

where the sum is intended over the Bravais lattice and \vec{k} and \vec{k}' belong to the reciprocal lattice.

(2 Points)

- The generic extended states of the graphene sheet have the form:
 $|\psi\rangle = \sum_{\vec{k}} c_{A\vec{k}} |\phi_{A\vec{k}}\rangle + c_{B\vec{k}} |\phi_{B\vec{k}}\rangle$, where $c_{A\vec{k}}$ and $c_{B\vec{k}}$ are complex numbers. Find the eigenstates of the system and the corresponding eigenvalues. How many bands do you find? What are the energy-wavevector relations? (2 Points)

Hint: You will need to solve a 2×2 matrix eigenvalue problem - the time-independent Schrödinger equation written in the basis of extended states.

- Show that the Fermi energy is equal to zero by verifying that this gives the correct electron density. Find the set of \vec{k} points for which $\epsilon(\vec{k}) = 0$. Show that these correspond to the corners of the first Brillouin zone. (2 Points)
- Plot the complete dispersion relation of graphene. (1 Point)
- (Oral)** Show that near to a (first) Brillouin zone corner with wavevector \vec{K} (note that \vec{K} is the location of a BZ corner, not a reciprocal lattice vector), the spectrum is approximately

$$\epsilon(\vec{k}) \approx \pm \hbar v \sqrt{(k_x - K_x)^2 + (k_y - K_y)^2}.$$

Determine the velocity v in terms of t and a_C . This behaviour is intermediate between that of a metal (with a Fermi surface instead of Fermi points \vec{K}) and an insulator (with a band gap). Indeed, graphene behaves like a semi-metal, with poor conductivity yet significantly higher than in an insulator.

Frohes Schaffen!